

Notes on the birds of Laquipampa Wildlife Refuge, Lambayeque, Peru

by Fernando Angulo Pratolongo, Jeremy N. M. Flanagan, Willem-Pier Vellinga & Napoleon Durand

Received 12 October 2011

SUMMARY.—Laquipampa was first designated a Reserved Zone, a temporary protected area category used by the Peruvian government until more complete studies permit a final designation, in 1982. The initial designation was based on the presence of the endemic and globally threatened White-winged Guan *Penelope albipennis*, but few other data on its other avifauna were available. We present the first comprehensive bird list (187 species) for this important reserve based on observations in 2001–10 and provide comments on several noteworthy records.

Laquipampa lies within the Incahuasi district of Ferreñafe province in Lambayeque. With the rediscovery of the presumed extinct White-winged Guan *Penelope albipennis* in Quebrada San Isidro in 1977 (see De Macedo 1979), further searches were initiated in Lambayeque and Piura. Ortiz (1980) found the species in two valleys at Laquipampa: Negrahuasi and Reloj. Based on this, a protected area was proposed and, in 1982, 11,346 ha were declared the Zona Reservada de Laquipampa to protect the guan (El Peruano 1982). However, apart from guan surveys (*cf.* Ortiz 1980, Ortiz & Díaz 1997), little was known of the area's avifauna and it received little government interest for almost two decades. In 1998 the Instituto Nacional de Recursos Naturales (INRENA, now the Servicio Nacional de Areas Naturales Protegidas, SERNANP) appointed the area's first refuge manager and in 2001 the first park ranger was assigned. In July 2006 the area was finally designated a Refugio de Vida Silvestre (El Peruano 2006) or wildlife refuge, covering 8,328 ha. Here we present the first comprehensive bird list for the conservation area, with notes on species of interest.

Study area and field work

Laquipampa Wildlife Refuge (LWR) is sited on a ridge on the north bank of the río La Leche (known as the Moyán above its confluence with the Sangana). Elevation in the refuge ranges from 200 m in the south-west (06°24'30"S, 79°33'45"W) to 2,600 m, the peak of 'La Punta' (06°15'55"S, 79°28'09"W), in the extreme north-east (Fig. 1).

According to CDC-UNALM (2006), the lower part of LWR belongs to the 'Piura and Tumbes Dry Forests' ecoregion and the upper part to the 'West Montane forests of the North Andes' ecoregion (Fig. 1). The dry forest is semi-deciduous and is found from 200 m to 1,300–1,400 m, and dominant species include *Loxopterygium huasango* (Anacardiaceae), *Bursera graveolens* (Burseraceae) and *Eriotheca ruizii* (Malvaceae). It has leaf cover in the wet season (December–May). Humidity and vegetation density increase with altitude. Montane forest is found in the upper LWR at 1,400–1,800 m, the vegetation does not shed its leaves in the dry season, and is considerably more humid than dry forest, reflected in the biomass of orchids, ferns and lichens. Another important floral community is *Sural* and scrub, which is dominated by *Suro* bamboo (*Chusquea* sp.: Poaceae) and many shrubs, with *Barnadesia* spp. (Asteraceae) dominant. This habitat occurs at 1,800–2,600 m and is characterised by low tree density, except on the sides of ravines. Agricultural areas also exist in the reserve including fields, mostly of corn, rice and sugarcane, near the ríos Moyán and La Leche, and formerly




Figure 1. Map showing the location of the Laquipampa Wildlife Refuge in north-west Peru, its extent and the location of the five sites mentioned in the text. Site 1: Quebrada Negrahuasi, c.590 m. Site 2: Quebrada Shambo, c.1,360 m. Site 3: Corral Grande in Quebrada Salsipuedes, c.1,830 m. Site 4: lower reaches, c.350 m. Site 5: El Porongo, c.2,100 m.

used areas on level terrain dominated by *Acacia macracantha* (Fabaceae) at all elevations (established prior to the creation of the protected area).

On 21–31 August 2001, in the dry season, an expedition by ProAves Perú and Asociación Cracidae Perú produced the first bird list of 110 species (Flanagan & Angulo 2003). Two locations were visited: Quebrada Negrahuasi (22–24 August, 590 m, site 1) and Quebrada Shambo (25–29 August, c.1,000 m–1,400 m, site 2). In February 2006, a second expedition by Asociación Cracidae Perú explored the area in the wet season. Three camps were established, in Quebrada Negrahuasi (12–14 February, 590 m, site 1), Quebrada Shambo (15–19 February, c.1,360 m, site 2) and Corral Grande in Quebrada Salsipuedes (20–24 February, 1,830 m, site 3). A total of 130 species, 37 of them ‘new’ to the area (Angulo & Alemán de Lama 2006), was identified. Most records described here were made during these two visits. Other visits were in the dry season: on 11–13 July 2008 (at c.350 m, site 4) by FAP & JNMF, and 11–14 November 2010 (c.2,100 m, site 5, El Porongo) by FAP. An additional source is the records made during a two-year migrant bird project (September 2006–July 2008) conducted by ECOAN. Also included are records made around Moyán town, just outside LWR but with similar habitats and elevation, on the road to Incahuasi in February 2006. Finally, documented (with photographs) reports by local people are also included.

Results

The Appendix presents an annotated species list for LWR totaling 187 species. Of these, seven are globally threatened—one Critically Endangered (CR) and six Vulnerable (VU) (BirdLife International 2012), 34 are restricted-range Tumbesian endemics and two are restricted-range Southern Central Andes endemics (Stattersfield *et al.* 1998). Sound-recordings have been archived at www.xeno-canto.org and are indicated by their catalogue number, e.g. XC66383. Noteworthy records not previously reported in the literature are detailed below.

COMB DUCK *Sarkidiornis melanotos*

Recorded twice in Laquipampa (5 June 2005 and 2 February 2006) at Puchaca (200–350 m), on both occasions singles in rice fields beside the río La Leche. The species is rather rare in north-west Peru, but was recorded during 1979–2010 at the Eten marshes near Chiclayo (Angulo *et al.* 2010). Approximately ten other records (usually of singles) in 2005–12 are available online (Cornell Laboratory of Ornithology & National Audubon Society 2011), from Rica Playa, in Tumbes, Laguna Ñapique (max. 25), Batanes and Puente Internacional, in Piura, and Olmos, Bosque de Pomac and Tinajones Reservoir (max. 55) in Lambayeque, the latter being the closest locality to Laquipampa. There is also a record from the río Tocto, Lambayeque (C. Hesse *in litt.* 2010).

BEARDED GUAN *Penelope barbata* VU

This restricted-range species was first reported at LWR in August 2001, in the Quebrada Shambo, where up to ten were seen at 1,300–1,700 m (Flanagan & Angulo 2002), and it was also heard in Quebrada Pescadera, a forested valley north of the refuge and above the village of Salas. In February 2006, *P. barbata* was found near the Shambo and Corral Grande camps, at 1,350–1,830 m. In November 2010 it was recorded to 2,150 m at El Porongo. On 17 February 2006, near Shambo camp at 1,400 m, a pair was found with a chick estimated on plumage to be *c.* 2 weeks old (Delacour & Amadon 2004), suggesting that breeding here coincides with the wet season (December–February). The estimated population at LWR is 20–25 individuals based on available habitat.

WHITE-WINGED GUAN *Penelope albipennis* CR

First recorded at LWR in 1978 in Quebrada Negrahuasi, where 5–6 individuals were seen (Ortiz 1980). In 1987, 2–4 were recorded there (Ortiz & Díaz 1997). In 2000–01, FAP surveyed Quebrada Negrahuasi and two other sites—Lajas and Quebrada Shambo—on the advice of local people, finding six, 11 and two guans respectively (Angulo 2000, 2001). As of 2005, the refuge supported a min. population of 22 individuals (Angulo 2008). At LWR, White-winged Guan has been found at 600–1,400 m and also in the same quebrada, Shambo, as Bearded Guan, where they share a narrow altitudinal band at 1,300–1,400 m. The species is recorded regularly at Lajas, just below the village of Laquipampa (outside the refuge), and is also known to descend at dawn and dusk to a place *c.* 500 m north of the village of Laquipampa, accessed via some fields. Moreover, ND noted that the species feeds in the village itself when a certain *Ficus* sp. (Moraceae) tree is fruiting. In 2007 eight captive-bred birds were reintroduced into Quebrada Negrahuasi (Angulo 2008) and had two chicks three months later. The estimated population in the refuge is 25–30 individuals.




Figure 2. Rufous-necked Wood Rail *Aramides axillaris*, Quebrada Shambo, Laquipampa Wildlife Refuge, Peru, 15 February 2007 (Oscar Rodríguez)

Figure 3. Captive Ruddy Quail-Dove *Geotrygon montana*, claimed to have been trapped locally, Laquipampa, Peru, 3 October 2005 (Napoleon Durand)

Figure 4. Mouse-coloured Tyrannulet *Phaeomyias murina tumbezana*, Laquipampa Wildlife Refuge, Peru, 22 February 2006 (Willem-Pier Vellinga)

Figure 5. Male Blue Seedeater *Amaurospiza concolor*, Corral Grande, Laquipampa Wildlife Refuge, Peru, 24 February 2006 (Willem-Pier Vellinga)

RUFIOUS-NECKED WOOD RAIL *Aramides axillaris*

Photographed by O. Rodríguez in Quebrada Shambo at 1,270 m on 15 February 2007 (Fig. 2). *A. axillaris* was first recorded in Peru in Tumbes Reserved Zone (now Cerros de Amotape National Park) in 1986 (Parker *et al.* 1995). In Tumbes it has been reported from mangroves (Valqui & Walker 2002) and dry forest near El Caucho and Pozo del Pato in the north of Cerros de Amotape National Park (Walker 2002). Our LWR record extends the species' distribution c.280 km south. According to (unconfirmed) information from the park rangers, the species is present at LWR year-round, but is more active and vocal in January–April.

OCHRE-BELLIED DOVE *Leptotila ochraceiventris* VU

Heard at 550–2,100 m, with one poor sound-recording (XC38631) obtained at Quebrada Shambo in February 2006, and it was recorded again at El Porongo in November 2010. These records appear to be the southernmost published. Further north it has been reported at Palambla (Koepcke 1961).

[RUDDY QUAIL-DOVE *Geotrygon montana*

One kept as a ‘pet’ by a local person (Fig. 3), who claimed that the bird was captured in the valley. Its origin is of course unclear. In Amazonia (and possibly elsewhere), the species reportedly undertakes irruptive movements (Stouffer & Bierregaard 1993), so our record could be related to the closest known west-slope population (c.300 km to the north in Loja, Ecuador (Ridgely & Greenfield 2001), or perhaps to that in western Amazonia.)

KOEPCKE’S SCREECH OWL *Megascops koepckeeae*

Sound-recorded near the Corral Grande camp in February 2006: during the night one was heard calling regularly and two birds, presumably male and female, were counter-singing just before dawn (XC5526, 8649). The species was also heard at night at 2,150 m at El Porongo. This is the northernmost record of the species on the west slope of the Andes. Several apparently well-documented records from the nearest localities further south (Sinsicap and Bosque Cachil) are unpublished (but see Cornell Laboratory of Ornithology & National Audubon Society 2011).

OILBIRD *Steatornis caripensis*

One observed on a tree was photographed by JNMF in Quebrada Shambo at c.1,400 m in August 2001. Schulenberg *et al.* (2007) stated ‘very rarely is reported from coast’. It has been recorded in Pimentel and Chiclayo (FAP unpubl.). The closest-known caves are apparently near San Andrés de Cutervo (SERNANP 2010) on the east slope of the Andes 77 km east of LWR.

LITTLE WOODSTAR *Chaetocercus bombus* VU

One photographed by A. Gonzales de la Cruz in Quebrada Shambo at 1,317 m, in May 2008, matches the species’ presumed habitat preference for humid and semi-deciduous forests. Little Woodstar was formerly considered threatened based on the paucity of late-20th century records (implying a decrease in numbers) and the destruction of humid forests (Collar *et al.* 1992). However, recent years have witnessed increasing records from the northern part of the historical range in Peru, especially the Marañón basin in Cajamarca, Piura, Amazonas and Lambayeque (Cornell Laboratory of Ornithology & National Audubon Society 2011). However, little is known of its ecology (Collar *et al.* 1992) and it would be interesting to clarify the species’ status in LWR.

RUFOUS-NECKED FOLIAGE-GLEANER *Syndactyla ruficollis* VU

In Peru known from the west slope of the northern Andes and from the Amotape range at 600–2,650 m (Schulenberg *et al.* 2007). After Cerros de Amotape National Park and El Angulo Hunting Reserve, LWR becomes the third protected area known to harbour the species in Peru and the only one in the main Andean chain. It was found at 1,360–2,150 m.

HENNA-HEADED FOLIAGE-GLEANER *Hylodyptes erythrocephalus* VU

In Peru protected areas known to support this species are Cerros de Amotape National Park and El Angulo Hunting Reserve, making LWR the third conservation unit in the country

and the first in the main Andean chain. In LWR it was encountered at Negrahuasi and Shambo, between c.550 and 2,100 m.

RUSTY-BREASTED ANTPITTA *Grallaricula ferrugineipectus leymebambae*

The song of *G. ferrugineipectus* was recorded at 2,150 m at El Porongo (XC66383). In recent years the species has been encountered at various locations on the west slope in Peru and Ecuador. In Ecuador, where it was previously unknown, the species has been found north to dpto. Pichincha, and in Peru it was initially found on the west slope near Canchaque (Schulenberg 1981) and subsequently near Ayabaca by P. Coopmans (Vellinga *et al.* 2004). Ours is the southernmost record (c.100 km south of Canchaque) and the first for Lambayeque.

MOUSE-COLOURED TYRANNULET *Phaeomyias murina*

A small confusing flycatcher that was trapped and observed several times in Quebrada Shambo (c.1,400 m) and Corral Grande (c.1,800 m) was identified as *P. m. tumbezana*. Our photographs compare well with specimens of *P. m. tumbezana* (Fig. 4) and the illustration in Schulenberg *et al.* (2007). In less-than-ideal field conditions the conspicuous wingbars and eyestripe, and the dark crown could cause confusion with Grey-breasted Flycatcher *Lathotriccus griseipectus*, rather than with *P. m. inflava*. On current knowledge, *P. m. inflava* occurs from the lowlands of southern Piura south, whereas *P. m. tumbezana* occurs in the lowlands of northern Piura north to Ecuador, but occurs south at least to the Olmos–Bagua highway, e.g. at 1,800 m at Limon de Porculla (D. F. Lane *in litt.* 2009). *P. m. inflava* has not yet been found in LWR but is known from Bosque de Pomac Historical Sanctuary, lower down in the floodplain of the río La Leche (06°29'S 79°46'W; 100–200 m). It is unclear if *P. m. inflava* and *P. m. tumbezana* co-occur in LWR. For now, there are no records at 500–1,400 m. Vocally, *P. m. inflava* and *P. m. tumbezana* are apparently similar (Schulenberg *et al.* 2007), although one vocalisation of *P. m. tumbezana* (XC38632) is perhaps unique, as we have not heard it from *P. m. inflava*.

GREY-BREASTED FLYCATCHER *Lathotriccus griseipectus* VU

Found at all three 2006 camps. It was sound-recorded in Quebrada Shambo (XC5626). LWR is south of the presumed range of the species, but it has recently been encountered even further south in the Zaña Valley as well (FAP unpubl.).

PIURA CHAT-TYRANT *Ochthoeca piurae* NT

This Peruvian endemic is known from Palambla and Porculla (Piura), Samne and Sinsicap (La Libertad), and Colcabamba, Wiñapatun, Noqno and San Damian (Ancash) (Collar *et al.* 1992). LWR is the first known location for Lambayeque and the first protected area known to harbour the species, when one was mist-netted near Corral Grande on 23 February 2006. At a similar elevation on the road to Incahuasi, just outside LWR (Moyán), the species was also observed and sound-recorded (XC5228) on 12 February 2006.

BLUE SEEDEATER *Amaurospiza concolor*

A male was trapped on 24 February 2006 near Corral Grande, at c.1,840 m in a mist-net within *Chusquea* sp. (Fig. 5). A presumed female and a male were observed by WPV at the same place one day earlier. LWR represents only the third-known location for the species in Peru; it was found in the Zaña Valley in 1996 (Salinas *et al.* 1998) and in a scrubby area west of Porculla pass at c.1,800 m (B. Walker *in litt.* 2009).

Discussion

The preliminary report on the 2001 expedition was instrumental in securing the present status of wildlife refuge, and the results presented here reinforce the importance of LWR for bird conservation in the Tumbesian Endemic Bird Area. The main importance of LWR remains the population of White-winged Guan; this is the only state-protected area known to date. Recently, two Regional Conservation Areas in which the species occurs have been created, in Piura and Lambayeque.

LWR also harbours a large number of Tumbesian endemics. Of the 55 range-restricted species of the Tumbesian region (*cf.* Stattersfield *et al.* 1998), 34 occur in LWR, making it meritorious of broader recognition as a key site in the protection of Tumbesian natural heritage. LWR is also crucial due to the altitudinal gradient of undisturbed forest protected therein, with species more associated with montane cloud forest present at higher altitudes, e.g. Bearded Guan. Such elevational transects of intact forest are increasingly scarce on the west slope of the northern Andes in Peru. Further field work could still discover other scarce species, e.g., Grey-headed Antbird *Myrmeciza griseiceps*, which is known both north and south of LWR in the main Andean chain. Such field work should focus on upper montane forests (2,000–2,600 m) and include seasons other than those during which we visited.

LWR lies immediately south of Comunidad Campesina Laquipampa and north of Salas. The economy of these communities mostly focuses on agriculture and cattle. Major threats to the refuge include its excessive use by cattle and land invasions by people.

Acknowledgements

Field work in 2001 was funded by the American Bird Conservancy and World Pheasant Association. 2006 field work was supported by BirdLife International via funds from the British Birdfair. Other visits were supported by Asociación Cracidae Perú. The people of Laquipampa are thanked for their friendship and assistance throughout, especially Simón Mendo, Simón Vásquez, Nazario Rojas, Marta Llaquento, Matias Santa Cruz, Wilmer de la Cruz and Winston de la Cruz. Team members at different times have included: Todd Mark, Giacomo Podesta, Carlos Benites, Pablo Venegas and Jefferson Diaz. Alex More is thanked for his help in the field and for producing the map. Dan Lane, Letty Salinas and Alfredo Begazo assisted with species identification from sound-recordings and photographs. Barry Walker shared his notes on Blue Seedeater. Constantino Auca permitted use of data from the ECOAN / NMBCA migratory bird study and its field workers Oscar Rodríguez, Daniel Diaz and Alejandro Gonzales. INRENA / SERNANP personnel Anibal Calderon, Dante Aleman, Vicente Cortez and Leticia Vasquez provided study permits.

References:

- Angulo, F. 2000. Informe de la salida de campo a la Zona Reservada de Laquipampa (29 febrero al 07 de marzo de 2000). Reporte interno 1. Asociación Cracidae Perú, Chiclayo.
- Angulo, F. 2001. Informe de la salida de campo a la Zona Reservada de Laquipampa (21 al 28 de agosto de 2001). Reporte interno 3. Asociación Cracidae Perú, Chiclayo.
- Angulo, F. 2008. Current status and conservation of wild and reintroduced White-winged Guan (*Penelope albipennis*). *Orn. Neotrop.* 19 (Suppl.): 279–286.
- Angulo, F. 2009. Peru. Pp. 307–316 in Devenish, C., Díaz Fernández, D. F., Clay, R. P., Davidson, I. & Yépez Zabala, I. (eds.) *Important Bird Areas Americas: priority sites for biodiversity conservation*. BirdLife International, Quito.
- Angulo, F. & Alemán de Lama, D. (eds.) 2006. Expedición Laquipampa 2006. Informe preliminar, marzo 2006. Asociación Cracidae Perú, Chiclayo.
- Angulo Pratonlongo, F., Tello, J. G. & Barrio, J. 2008. First records of Fasciated Tiger Heron *Tigrisoma fasciatum* in the west Peruvian Andes. *Cotinga* 29: 173–175.
- Angulo P., F., Schulenberg, T. S. & Puse F., E. 2010. Las aves de los humedales de Eten, Lambayeque, Perú. *Ecol. Aplicada* 9: 71–81.
- BirdLife International. 2012. IUCN Red List for birds. www.birdlife.org (accessed 10 May 2012).
- Centro de Datos para la Conservación–Universidad Nacional Agraria La Molina (CDC-UNALM). 2006. *Análisis de la cobertura ecológica del Sistema Nacional de Áreas Naturales Protegidas por el estado*. CDC-UNALM/TNC, Lima.

- Collar, N. J., Gonzaga, L. P., Krabbe, N., Madroño Nieto, A., Naranjo, L. G., Parker, T. A. & Wege, D. C. 1992. *Threatened birds of the Americas: the ICBP / IUCN Red Data book*. International Council for Bird Preservation, Cambridge, UK.
- Cornell Laboratory of Ornithology & National Audubon Society. 2011. eBird: an online database of bird distribution and abundance. www.avianknowledge.net (accessed 27 February 2012).
- Delacour, J. & Amadon, D. 2004. *Curassows and related birds*. Second edn. Lynx Edicions, Barcelona & Amer. Mus. Nat. Hist., New York.
- De Macedo, H. 1979. Extinct bird found in Peru. *Oryx* 15: 33–37.
- El Peruano. 1982. Resolución Ministerial 00692 – AG-DGFF del 5 de octubre de 1982. Establecen la Zona Reservada Laquipampa para la preservación de la Pava Aliblanca.
- El Peruano. 2004. Decreto Supremo 034 – AG del 22 de septiembre de 2004. Categorización de especies amenazadas de fauna silvestre.
- El Peruano. 2006. Decreto Supremo 045 – AG del 07 de julio 2006. Categorizar de forma definitiva la Zona Reservada Laquipampa como “Refugio de Vida Silvestre Laquipampa”, ubicado en el departamento de Lambayeque.
- Flanagan, J. N. M. & Angulo, F. 2002. The White-winged Guan (*Penelope albipennis*), Bearded Guan (*P. barbata*) and the Zona Reservada de Laquipampa, NW Peru. *Bull. Cracid Specialist Group* 15: 27–31.
- Flanagan, J. N. M. & Angulo, F. 2003. La Zona Reservada de Laquipampa – ecología, conservación y manejo. *In* Memorias del Primer Congreso Internacional de Bosques Secos (6–9 Nov. 2003), Univ. de Piura.
- Koepcke, M. 1961. Birds of the western slope of the Andes of Peru. *Amer. Mus. Novit.* 2028: 1–31.
- Ortíz, T., E. 1980. Estudio preliminar sobre la “Pava de Ala Blanca” (*Penelope albipennis*). Unpubl. rep. to Dirección General Forestal y de Fauna, Ministerio de Agricultura, Lima.
- Ortíz T., E. & Díaz, V. R. 1997. Estudio de campo y reevaluación de la población de Pava Aliblanca (*Penelope albipennis*). Pp. 218–232 in Strahl, S. D., Beaujon, S., Brooks, D. M., Begazo, A. J., Sedaghatkish, G. & Olmos, F. (eds.) *The Cracidae: their biology and conservation*. Hancock House, Blaine, WA.
- Parker, T. A., Schulenberg, T., Kessler, M. & Wust, W. H. 1995. History and conservation of the endemic avifauna in north-west Peru. *Bird Conserv. Intern.* 5: 201–233.
- Ridgely, R. S. & Greenfield, P. J. 2001. *The birds of Ecuador*. Cornell Univ. Press, Ithaca, NY.
- Salinas, L., Franke, I., Samamé, M. & Fjeldså, J. 1998. Primer registro de *Amaurospiza concolor* (Emberizidae, Passeriformes) para el Perú. *Resúm. VII Reunión Cient. ICBAR*: 107.
- Schulenberg, T. S. & Parker, T. A. 1981. Status and distribution of some northwest Peruvian birds. *Condor* 83: 209–216.
- Schulenberg, T. S., Stotz, D. F., Lane, D. F., O’Neill, J. P. & Parker, T. A. 2010. *Birds of Peru*. Princeton Univ. Press
- Servicio Nacional de Áreas Naturales Protegidas (SERNANP). 2010. *Plan maestro del Parque Nacional Cutervo 2010–2014*. SERNANP, Lima.
- Stattersfield, A. J., Crosby, M. J., Long, A. J. & Wege, D. C. 1998. *Endemic Bird Areas of the world: priorities for biodiversity conservation*. BirdLife International, Cambridge, UK.
- Stouffer, P. C. & Bierregaard, R. O. B. 1993. Spatial and temporal abundance patterns of Ruddy Quail-Doves (*Geotrygon montana*) near Manaus, Brazil. *Condor* 95: 896–903.
- Valqui, T. & Walker, B. 2002. Importance of mangrove forests in Peru with notes on Bare-throated Tigerheron *Tigrisoma mexicanum* and Rufous-necked Wood-rail *Aramides axillaris*. *Cotinga* 18: 58–61.
- Vellinga, W. P., Flanagan, J. N. M. & Mark, T. R. 2004. New and interesting records of birds from Ayabaca province, Piura, north-west Peru. *Bull. Brit. Orn. Cl.* 124: 124–142.
- Walker, B. 2002. Observations from the Tumbes Reserved Zone, dpto. Tumbes, with notes on some new taxa for Peru and a checklist of the area. *Cotinga* 18: 37–43.

Addresses: Fernando Angulo Pratonlongo, Asociación Cracidae Perú & CORBIDI. Jeremy N. M. Flanagan, Asociación Tu Tierra, Chiclayo, Peru. Willem-Pier Vellinga (corresponding author), Stichting xeno-canto voor natuurgeluiden (xeno-canto foundation), Groenesteinlaan 12, 9722 BX, Groningen, the Netherlands, e-mail: wp@xeno-canto.org. Napoleon Durand, Refugio de Vida Silvestre Laquipampa, Servicio Nacional de Áreas Naturales Protegidas por el Estado (SERNANP), Av. Víctor Muro 1175, Ferreñafe, Lambayeque, Peru.

APPENDIX: birds of Laquipampa Wildlife Refuge

X = recorded in the season indicated. An empty box indicates a lack of records at the relevant season, and - = a degree of certainty that the species was not present. Evidence: C = mist-netted, photographed and released, P = photographed, R = sound-recorded, V = seen only, H = heard only, O = just outside LWR limits. Photographs include species photographed in the wild or birds photographed after being captured in the area by local people. Also shown are species indicative of Endemic Bird Areas; T = Tumbesian EBA, SCA = South Central Andes EBA, or country endemics (P) and threat status according to both IUCN (BirdLife International 2012) and the Peruvian government (El Peruano 2004).

	Elevation min. max.	Season		Evidence	Threat IUCN Peru	Endemic EBA Peru	Remarks
		wet	dry				
<i>Ixobrychus exilis</i>	250 250	X	X	V			
<i>Nycticorax nycticorax</i>	550 550	X	X	V			
<i>Butorides striata</i>	300 550	X	X	V			
<i>Bubulcus ibis</i>	300 800	X	X	V			
<i>Ardea alba</i>	300 550	X	X	P			
<i>Egretta thula</i>	300 1,100	X	X	P			
<i>Egretta caerulea</i>	300 400	X	X	P			
<i>Cathartes aura</i>	300 2,100	X	X	P			
<i>Coniopsis atratus</i>	300 1,830	X	X	V			
<i>Sarcorampus papa</i>	300 1,700	X	X	V			
<i>Vultur gryphus</i>	2,200 2,600	X	X	V	EN		
<i>Gampsonyx swainsonii</i>	550 550	X	X	P			
<i>Accipiter striatus</i>	1,830 2,100	X	X	V			
<i>Accipiter bicolor</i>	300 1,500	X	X	V			
<i>Buteogallus urubitinga</i>	590 1,400	X	X	V			
<i>Buteogallus meridionalis</i>	200 400	X	X	P			
<i>Harpophalium solitarius</i>	800 800	X	X	V			
<i>Geranoetus melanoleucus</i>	300 2,100	X	X	V			
<i>Parabuteo unicinctus</i>	550 700	X	X	V			
<i>Buteo leucorhous</i>	2,000 2,100	X	X	V			
<i>Buteo polyosoma</i>	550 700	X	X	V			
<i>Caracara cheriway</i>	300 700	X	X	V			
<i>Falco sparverius</i>	300 800	X	X	P			
<i>Falco peregrinus</i>	300 1,400	X	X	V			
<i>Aramides axillaris</i>	1,360 1,360	X	X	P			
<i>Charadrius vociferus</i>	300 550	X	X	V			
<i>Burhinus superciliosus</i>	300 300	X	X	V			
<i>Actitis macularia</i>	550 550	X	X	V			
<i>Columbina cruziana</i>	300 1,500	X	X	P, R			XC21773
<i>Columba litua</i>	300 800	X	X	V			
<i>Patagioena fasciata</i>	1,360 2,150	X	X	P, R			XC69831
<i>Zenaidura macroura</i>	300 1,500	X	X	P, R			XC8661, 16719, 21777, 21779
<i>Zenaidura macroura</i>	300 2,150	X	X	P			
<i>Leptotilia verreauxi</i>	300 2,150	X	X	P, R			XC16720, 21594
<i>Leptotilia ochraceiventris</i>	550 2,100	X	X	R	VU	T	XC38632
<i>Aratinga wagleri</i>	590 1,830	X	X	V			
<i>Aratinga erythrogenys</i>	300 2,100	X	X	P, R		T	XC16721, 16724

	Elevation		Season		Evidence	Threat		Endemic		Remarks
	min.	max.	wet	dry		IUCN	Peru	EBA	Peru	
<i>Forpus coelestis</i>	300	1,800	X	X	P					
<i>Coccyzus erythrophthalmus</i>	700	700	X		P					
<i>Crotophaga sulcirostris</i>	300	1,800	X	X	R					XC21774
<i>Tapera naevia</i>	300	300	X	X	H					
<i>Tyto alba</i>	800	800	X	X	P					
<i>Megascops roboratus</i>	590	2,100	X	X	R					XC5627, 8660
<i>Megascops hoopceae</i>	1,830	2,150	X	X	R			P		XC5526, 8649
<i>Pulsatrix perspicillata</i>	600	700	X	X	H					
<i>Ciccaba albitarsis</i>	1,400	1,400	X	X	R					XC29018
<i>Glaucidium peruanum</i>	300	1,800	X	X	R					XC5609, 8644, 21592
<i>Athene cunicularia</i>	300	800	X	X	P					
<i>Stentornis caripensis</i>	1,400	1,400	X	X	P					
<i>Nyctibius griseus</i>	590	1,360	X	X	P, R					
<i>Chordeiles acutipennis</i>	300	1,400	X	X	P					
<i>Nyctidroma albicollis</i>	1,360	1,800	X	X	R					
<i>Caprimulgus anthomyi</i>	550	550	X	X	H					XC29018, 8645
<i>Streptoprocne rutilla</i>	590	1,360	X	X	P					
<i>Streptoprocne zonaris</i>	1,400	1,400	X	X	V					
<i>Chaetura brachyura</i>	1,830	1,830	X	X	V					
<i>Aeronautes montivagus</i>	590	2,100	X	X	R					XC5180
<i>Phaethornis griseogularis</i>	550	2,100	X	X	V					
<i>Colibri coruscans</i>	1,400	1,800	X	X	V					
<i>Adelomyia melanogenys</i>	550	2,150	X	X	R					XC5590
<i>Lesbia nuna</i>	1,830	2,100	X	X	C					
<i>Aglaeactis cupripennis</i>	2,100	2,150	X	X	V					
<i>Coeligena iris</i>	1,500	2,100	X	X	V					SCA
<i>Patagona gigas</i>	1,800	1,800	X	X	V					
<i>Helminthaster longirostris</i>	590	800	X	X	V					
<i>Myrtis fanny</i>	1,800	1,830	X	X	V					T
<i>Chaetocercus bombus</i>	1,300	1,300	X	X	V					T
<i>Myrmia micrura</i>	300	1,050	X	X	V					T
<i>Leucippus</i> sp.	590	590	X	X	V					T
<i>Amazilia amazilia</i>	300	1,830	X	X	C, R					XC5608
<i>Trogon mesurus</i>	550	1,800	X	X	P, R					XC5255-56
<i>Megaceryle torquata</i>	300	700	X	X	P					
<i>Chloroceryle americana</i>	300	700	X	X	V					
<i>Momotus momota</i>	590	590	X	X	P, R					XC5237

	Elevation		Season		Evidence	Threat	Endemic		Remarks
	min.	max.	wet	dry			EBA	Peru	
<i>Picumnus sclateri</i>	1,360	1,830	X	X	C, R		T		XC8633
<i>Picoides fumigatus</i>	1,830	1,920	X	X	C				
<i>Vermilionis callonotus</i>	300	1,360	X	X	V				
<i>Colaptes rubiginosus</i>	300	1,700	X	X	P, R				XC5244
<i>Dryocopus lineatus</i>	590	1,360	X	X	V				
<i>Campephilus gayaquilensis</i>	550	700	X	X	P				
<i>Furnarius leucopus</i>	300	1,830	X	X	P				
<i>Synallaxis azarae</i>	1,400	2,100	X	X	R				XC44728, 44730, 44732
<i>C-ratiolauca antisiensis</i>	1,800	2,150	X	X	C, R				XC5597
<i>Syndactyla ruficollis</i>	1,360	2,150	X	X	C, R	VU	T		XC5177, 8750
<i>Hylodyptilus erythrocephalus</i>	550	2,100	X	X	R	VU	T		XC8727
<i>Lepidocolaptes souleyetii</i>	300	1,400	X	X	P, R				XC8699
<i>Thamnophilus zurumae</i>	1,400	2,150	X	X	P, R				XC5181
<i>Thamnophilus bernardi</i>	300	1,650	X	X	P, R		T		XC5566, 21586
<i>Grallaria guatemalensis</i>	1,400	1,830	X	X	C, R		T		XC8747
<i>Grallaria ruficapilla</i>	1,400	2,150	X	X	R				many, e.g., XC8603
<i>Grallaria waikiki</i>	700	1,200	X	X	V		T		
<i>Grallaria ferruginepectus</i>	2,100	2,150	X	X	R				XC66383, 69835
<i>Scytalopus latrans subcinereus</i>	1,400	2,150	X	X	R				XC41723, 44728, 44732
<i>Melanopareia elegans</i>	300	1,830	X	X	C, R		T		XC5595, 8579
<i>Myiopigis subplacens</i>	700	1,350	X	X	R		T		XC5263
<i>Camptostoma obsoletum</i>	300	2,150	X	X	R				e.g. XC5179
<i>Mecocerculus calopterus</i>	1,830	1,830	X	X	C, R				XC8603
<i>Phaenomyia murina tumbezana</i>	1,400	2,150	X	X	C, R				XC38632
<i>Euscarthmus meloryphus</i>	300	2,150	X	X	R				XC5615, 16726
<i>Pseudelaenia leucospodia</i>	300	700	X	X	V		T		
<i>Mionectes striatocollis</i>	1,830	1,830	X	X	C				
<i>Todirostrum cinereum</i>	550	1,400	X	X	V				
<i>Myiophobus fuscatus</i>	550	1,350	X	X	V				
<i>Lathrotriccus griseipectus</i>	1,350	1,830	X	X	R	VU	T		XC5614
<i>Contopus cinereus</i>	590	1,830	X	X	P, R				XC5626
<i>Contopus junigatus</i>	1,360	1,830	X	X	R				XC5584
<i>Sayornis nigricans</i>	550	1,500	X	X	P				XC8731
<i>Pyrocephalus rubinus</i>	300	1,830	X	X	P				
<i>Myiotheretes striatocollis</i>	1,800	2,150	X	X	V				
<i>Tumbezia salvini</i>	590	590	X	X	P		T		
<i>Oothoeca piurae</i>	1,800	1,830	X	X	C, R	EN	T		XC5228

	Elevation min.	max.	Season		Evidence	Threat		Endemic		Remarks
			wet	dry		IUCN	Peru	EBA	Peru	
<i>Muscivalla brevicauda</i>	300	590	X	X	V					
<i>Baird's Flycatcher</i>	300	1,400	X	X	P, R		T			XC5236
<i>Streaked Flycatcher</i>	590	590	X		V					
<i>Snowy-throated Kingbird</i>	300	300	X	X	P					
<i>Tropical Kingbird</i>	300	1,830	X	X	R					
<i>Dusky-capped Flycatcher</i>	550	2,100	X	X	V, R					
<i>Sooty-crowned Flycatcher</i>	1,700	1,700	X	X	V		T			XC5238 e.g. XC8731
<i>Black-and-white Becard</i>	1,360	1,360	X		C					
<i>Rufous-browed Peppershrike</i>	300	2,150	X	X	P, R					XC5619
<i>White-tailed Jay</i>	300	1,830	X	X	P, R, C					XC5183, 5185
<i>Blue-and-white Swallow</i>	300	2,100	X	X	V					
<i>Southern Rough-winged Swallow</i>	590	590	X	X	V					
<i>Grey-breasted Martin</i>	300	1,400	X	X	V					
<i>Barn Swallow</i>	550	550	X		V					
<i>Chestnut-collared Swallow</i>	1,050	1,360	X	X	V					
<i>House Wren</i>	300	1,900	X	X	C, R					XC5599
<i>Fasciated Wren</i>	300	1,830	X	X	R					XC8581
<i>Speckle-breasted Wren</i>	1,150	1,700	X	X	V					
<i>Superciliated Wren</i>	300	1,830	X	X	R					
<i>Tropical Gnatcatcher</i>	300	1,500	X	X	R		T			XC5589, 21775
<i>Slaty-backed Nightingale-Thrush</i>	1,700	2,100	X	X	R					XC16725
<i>Plumbeous-backed Thrush</i>	300	1,830	X	X	P, R, C					XC41481
<i>Andean Slaty Thrush</i>	1,400	1,830	X	X	P, R, C					XC5567
<i>Great Thrush</i>	1,800	2,150	X	-	V, R					XC5176, 8732
<i>Chiguanco Thrush</i>	1,800	2,000	X	X	R, O					XC69842-43
<i>Long-tailed Mockingbird</i>	300	1,830	X	X	P, R					XC8700
<i>Parámo Pipit</i>	1,800	1,800	X	X	V					XC5604
<i>Black-and-white Tanager</i>	590	1,830	X	-	R					XC5603
<i>Black-eared Hemispingus</i>	1,900	2,150	X	X	R					XC6984-41
<i>Rufous-chested Tanager</i>	1,830	1,830	X		R					XC5596
<i>Blue-grey Tanager</i>	300	1,800	X	X	R					XC5558
<i>Blue-and-yellow Tanager</i>	1,800	1,800	X	X	V					
<i>Fawn-breasted Tanager</i>	1,830	1,830	X	X	V					
<i>Silvery Tanager</i>	1,360	1,830	X	X	V					
<i>Cinereous Conebill</i>	1,800	1,830	X	X	V					
<i>Rusty Flowerpiercer</i>	1,800	1,830	X	X	C					
<i>Plushcap</i>	2,100	2,100	X	X	V					

	Elevation		Season	Evidence	Threat	Endemic		Remarks
	min.	max.				EBA	Peru	
<i>Coereba flaveola</i>	300	1,500	wet X dry X	V				
<i>Saltator nigriceps</i>	1,800	2,150	X	P, R, C	T			XC5184
<i>Saltator striatipectus</i>	1,050	1,050	X	V				
<i>Zonotrichia capensis</i>	1,400	2,100	X	V				
<i>Rhyncospiza stolzmanni</i>	300	1,400	X	P, R	T			XC21781
<i>Phrygilus plebejus</i>	1,800	2,100	X	V				
<i>Piezorhina cinerea</i>	300	300	X	V	T		P	XC33256
<i>Poospiza hispaniolensis</i>	300	300	X	R				
<i>Sticis flaveola</i>	300	1,830	X	P				XC5582
<i>Volatinia jacarina</i>	300	1,830	X	R				
<i>Sporophila peruviana</i>	300	700	X	V				
<i>Sporophila telasco</i>	300	590	X	V				
<i>Arremon abeliei</i>	590	1,500	X	P, C	T			XC5169
<i>Arremon assimilis</i>	1,360	2,150	X	C, R				XC5173
<i>Atlapetes leucopterus</i>	600	1,830	X	C, R				
<i>Atlapetes albiceps</i>	300	2,150	X	V	T			
<i>Atlapetes seebohmi</i>	1,650	1,650	X	V	T			
<i>Piranga flava</i>	300	1,830	X	P, R				XC8580
<i>Pheucticus chrysogaster</i>	300	1,830	X	P, R, C				XC5182
<i>Amauroscoiza concolor</i>	1,830	1,830	X	C				
<i>Parula pitiayumi</i>	300	1,830	X	P, R				XC5174
<i>Geothlypis aquinoctialis</i>	1,400	1,830	X	R				XC5581, 5583
<i>Mnioborus miniatus</i>	1,360	2,100	X	C				
<i>Basilentus fraseri</i>	550	1,700	X	C, R	T			XC5178
<i>Basilentus nigrocrissatus</i>	1,830	2,100	X	C, R				XC8764
<i>Basilentus trifasciatus</i>	590	2,150	X	C, R	T			XC5187, 8648
<i>Icterus graccarinnae</i>	300	1,800	X	P, R	T			XC16723
<i>Icterus mesomelas</i>	590	1,360	X	P, R, C				XC8735
<i>Dives warszewitzi</i>	300	1,150	X	V				
<i>Molothrus bonariensis</i>	300	300	X	P				
<i>Sturnella heliaca</i>	300	1,500	X	R				XC21580
<i>Carduelis magellanica</i>	1,800	2,100	X	R				XC5600
<i>Carduelis psaltria</i>	1,800	1,800	X	V				
<i>Euphonia laniirostris</i>	300	2,100	X	P, R				XC5245